

Renormalizable minimal SO(10) GUT in 4D and 5D¹

Takeshi FUKUYAMA

Department of Physics, Ritsumeikan University, Kusatsu, Shiga, 525-8577 Japan

Abstract. This report is a review of the present status of GUT, especially renormalizable minimal SO(10) GUT, and its future prospect. It consists of two parts. In part I, I review how the minimal renormalizable supersymmetric SO(10) GUT, an SO(10) framework with only one **10** and one **126** Higgs multiplets in the Yukawa sector, is attractive because of its high predictivity. Indeed it not only gave a consistent predictions on neutrino oscillation data but also did reasonable and interesting values for Leptogenesis, LFV, muon g-2, neutrinoless double beta decay etc. However, this model suffers from problems, apart from the small deviations from the observed values, related to running of gauge couplings and proton decay. The gauge coupling unification may be spoiled due to the presence of intermediate scales much lighter than the grand unification (GUT) scale. In addition, the gauge couplings blow up around the GUT scale because of the presence of Higgs multiplets of large representations. In order to remedy these pathologies, in part II, we extend GUT into 5D. We propose two approaches: one is to consider the warped extra dimension, using the bulk Higgs profile to explain the intermediate energy scales. Another is to use the orbifold GUT. Both approaches are complementary to each other.

Keywords: GUT, extra dimension

PACS: 12.10.Dm, 04.50.+h, 12.60.Jv, 14.60.Pq

PART I –GUT IN 4D

SUSY GUT is the most promising candidate beyond the Standard Model (SM) [2]. SM is the very powerful theory but it has the application limit like the other great theories. There are discrepancies with observations like neutrino mass [3], muon g-2,... It also has the unsatisfactory points which strongly suggests more comprehensive theory:

SM has so many parameters. It does not explain quark-lepton mass spectra, mixing angles, all phases, Higgs mass stability against quantum correction, three different gauge couplings and their unification, and Dark Matters (DM) etc. Here we consider the theory beyond SM from bottom up approach. Top down approach from string theory is also interesting [4][5]. On these problems there are many approaches but it seems to be SUSY GUT which may solve the whole problems mentioned above. Of course there are still room to accept non SUSY GUT and even non GUT.

Even if we accept it, there are so many SUSY GUTs. What is the gauge group, SU(5), SO(10), E_6 [6], $E_8 \times E_8$ [4] ? So we need the other criterion to select the gauge symmetry. Anomaly free condition may be good candidate for it. Chiral symmetry must be preserved under quantum correction. SO(10) is the smallest group which is free from anomaly. Such an anomaly free condition has well meaning if the theory is renormalizable. Let us consider the structure of the existing theories. They have the form of

$$L = L_{ren} + \frac{L_1}{\Lambda_1}. \quad (1)$$

Here the first L_{ren} denotes renormalizable Lagrangian and the second unrenormalizable effective Lagrangian. For $SU(3) \times U(1)$ theory, L_1 implies the Fermi coupling $G_W J^\mu J_\mu$. For $SU(3) \times SU(2) \times U(1)$ SM, this L_1 becomes the renormalizable $g_2 J^\mu W_\mu$ term but new effective term appears in the seesaw mechanism,

$$L = L'_{ren} + Y_V^T \frac{1}{M_R} Y_V (LH)^2 \equiv L_{ren} + \frac{L_2}{\Lambda_2}. \quad (2)$$

Here $L'_{ren} \supset L'_{ren}$ and $\Lambda_2 (= O(10^{13} \text{GeV})) \gg \Lambda_1$. Thus the theory is expressed as the sum of renormalizable theory plus cut off effective action, and renormalizable Lagrangian becomes more involved as the energy scale goes high. In the limit of $\Lambda_2 = \infty$ SM is renormalizable. Thus it may be reasonable to consider the renormalizability as the guiding principle for model building beyond SM.

¹ This is a talk in the Workshop on GUT held at Ritsumeikan Univ. on Dec.17-19 2007 [1]

The group theoretical properties give strong constraints on quark-lepton of the same family but very weak on those of different families. As for family symmetry, see [7].

The successful gauge coupling unification in the minimal supersymmetric standard model (MSSM), strongly support the emergence of a supersymmetric (SUSY) GUT around $M_{\text{GUT}} \simeq 2 \times 10^{16}$ GeV. SO(10) is the smallest simple gauge group under which the entire SM matter content of each generation is unified into a single anomaly-free irreducible representation, **16** representation. This **16** representation includes right-handed neutrino and SO(10) GUT incorporates the see-saw mechanism [8]. Among several models based on the gauge group SO(10), the renormalizable minimal SO(10) model has been paid a particular attention, where two Higgs multiplets $\{\mathbf{10} \oplus \overline{\mathbf{126}}\}$ are utilized for the Yukawa couplings with matters $\mathbf{16}_i$ (i = generation). A remarkable feature of the model is its high predictivity of the neutrino oscillation parameters as well as reproducing charged fermion masses and mixing angles.

Minimal supersymmetric SO(10) model

First we give a brief review of the renormalizable minimal SUSY SO(10) model.^{2 3}.

Yukawa coupling. This model was first applied to neutrino oscillation in [10]. However it did not reproduce the large mixing angles. It has been pointed out that CP-phases in the Yukawa sector play an important role to reproduce the neutrino oscillation data [11]. More detailed analysis incorporating the renormalization group (RG) effects in the context of MSSM has explicitly shown that the model is consistent with the neutrino oscillation data at that time and became a realistic model [12]. We give a brief review of this renormalizable minimal SUSY SO(10) model. Yukawa coupling is given by

$$W_Y = Y_{10}^{ij} \mathbf{16}_i H_{10} \mathbf{16}_j + Y_{126}^{ij} \mathbf{16}_i H_{126} \mathbf{16}_j, \quad (3)$$

where $\mathbf{16}_i$ is the matter multiplet of the i -th generation, H_{10} and H_{126} are the Higgs multiplet of **10** and $\overline{\mathbf{126}}$ representations under SO(10), respectively. Note that, by virtue of the gauge symmetry, the Yukawa couplings, Y_{10} and Y_{126} , are, in general, complex symmetric 3×3 matrices. After the symmetry breaking pattern of SO(10) to $SU(3)_c \times SU(2)_L \times U(1)_Y$ via $SU(4)_c \times SU(2)_L \times SU(2)_R$ or $SU(5) \times U(1)$, we find that two pair of Higgs doublets in the same representation appear as the pair in the MSSM. One pair comes from $(\mathbf{1}, \mathbf{2}, \mathbf{2}) \subset \mathbf{10}$ and the other comes from $(\overline{\mathbf{15}}, \mathbf{2}, \mathbf{2}) \subset \overline{\mathbf{126}}$. Using these two pairs of the Higgs doublets, the Yukawa couplings of Eq. (3) are rewritten as

$$\begin{aligned} W_Y = & (U^c)_i \left(Y_{10}^{ij} H_{10}^u + Y_{126}^{ij} H_{126}^u \right) Q_j + (D^c)_i \left(Y_{10}^{ij} H_{10}^d + Y_{126}^{ij} H_{126}^d \right) Q_j \\ & + (N^c)_i \left(Y_{10}^{ij} H_{10}^u - 3Y_{126}^{ij} H_{126}^u \right) L_j + (E^c)_i \left(Y_{10}^{ij} H_{10}^d - 3Y_{126}^{ij} H_{126}^d \right) L_j \\ & + L_i \left(Y_{126}^{ij} \nu_T \right) L_j + (N^c)_i \left(Y_{126}^{ij} \nu_R \right) (N^c)_j, \end{aligned} \quad (4)$$

where u_R , d_R , ν_R and e_R are the right-handed SU(2)_L singlet quark and lepton superfields, q and ℓ are the left-handed SU(2)_L doublet quark and lepton superfields, $H_{10}^{u,d}$ and $H_{126}^{u,d}$ are up-type and down-type Higgs doublet superfields originated from H_{10} and H_{126} , respectively, and the last term is the Majorana mass term of the right-handed neutrinos developed by the VEV of the $(\overline{\mathbf{10}}, \mathbf{1}, \mathbf{3})$ Higgs, ν_R . The factor -3 in the lepton sector is the Clebsch-Gordan coefficient.

In order to preserve the successful gauge coupling unification, suppose that one pair of Higgs doublets given by a linear combination $H_{10}^{u,d}$ and $H_{126}^{u,d}$ is light while the other pair is heavy ($\geq M_{\text{GUT}}$). The light Higgs doublets are identified as the MSSM Higgs doublets (H_u and H_d) and given by

$$\begin{aligned} H_u &= \tilde{\alpha}_u H_{10}^u + \tilde{\beta}_u H_{126}^u, \\ H_d &= \tilde{\alpha}_d H_{10}^d + \tilde{\beta}_d H_{126}^d, \end{aligned} \quad (5)$$

where $\tilde{\alpha}_{u,d}$ and $\tilde{\beta}_{u,d}$ denote elements of the unitary matrix which rotate the flavor basis in the original model into the (SUSY) mass eigenstates (See (43) in detail). Omitting the heavy Higgs mass eigenstates, the low energy superpotential

² This part is based on the works by T. Fukuyama, A. Ilakovac, T. Kikuchi, S. Meljanac and N. Okada

³ There is another flow of non-renormalizable minimal SO(10) GUT [9]

TABLE 1. The input values of $\tan\beta$, $m_s(M_Z)$ and δ in the CKM matrix and the outputs for the neutrino oscillation parameters.

$\tan\beta$	$m_s(M_Z)$	δ	σ	$\sin^2 2\theta_{12}$	$\sin^2 2\theta_{23}$	$\sin^2 2\theta_{13}$	$\Delta m_{\odot}^2/\Delta m_{\oplus}^2$
40	0.0718	93.6°	3.190	0.738	0.900	0.163	0.205
45	0.0729	86.4°	3.198	0.723	0.895	0.164	0.188
50	0.0747	77.4°	3.200	0.683	0.901	0.164	0.200
55	0.0800	57.6°	3.201	0.638	0.878	0.152	0.198

is described by only the light Higgs doublets H_u and H_d such that

$$\begin{aligned}
W_Y &= (U^c)_i \left(\alpha^u Y_{10}^{ij} + \beta^u Y_{126}^{ij} \right) H_u Q_j + (D^c)_i \left(\alpha^d Y_{10}^{ij} + \beta^d Y_{126}^{ij} \right) H_d Q_j \\
&+ (N^c)_i \left(\alpha^u Y_{10}^{ij} - 3\beta^u Y_{126}^{ij} \right) H_u L_j + (E^c)_i \left(\alpha^d Y_{10}^{ij} - 3\beta^d Y_{126}^{ij} \right) H_d L_j \\
&+ L_i \left(Y_{126}^{ij} v_T \right) L_j + (N^c)_i \left(Y_{126}^{ij} v_R \right) (N^c)_j,
\end{aligned} \tag{6}$$

where the formulas of the inverse unitary transformation of Eq. (5), $H_{10}^{u,d} = \alpha^{u,d} H_{u,d} + \dots$ and $H_{126}^{u,d} = \beta^{u,d} H_{u,d} + \dots$, have been used. Note that the elements of the unitary matrix, $\alpha^{u,d}$ and $\beta^{u,d}$, are in general complex parameters, through which CP-violating phases are introduced into the fermion mass matrices.

Providing the Higgs VEVs, $H_u = v \sin\beta$ and $H_d = v \cos\beta$ with $v = 174\text{GeV}$, the quark and lepton mass matrices can be read off as

$$\begin{aligned}
M_u &= c_{10} M_{10} + c_{126} M_{126} \\
M_d &= M_{10} + M_{126} \\
M_D &= c_{10} M_{10} - 3c_{126} M_{126} \\
M_e &= M_{10} - 3M_{126} \\
M_T &= c_T M_{126} \\
M_R &= c_R M_{126},
\end{aligned} \tag{7}$$

where M_u , M_d , M_D , M_e , M_T , and M_R denote the up-type quark, down-type quark, Dirac neutrino, charged-lepton, left-handed Majorana, and right-handed Majorana neutrino mass matrices, respectively. Note that all the quark and lepton mass matrices are characterized by only two basic mass matrices, M_{10} and M_{126} , and four complex coefficients c_{10} , c_{126} , c_T and c_R , which are defined as $M_{10} = Y_{10} \alpha^d v \cos\beta$, $M_{126} = Y_{126} \beta^d v \cos\beta$, $c_{10} = (\alpha^u/\alpha^d) \tan\beta$, $c_{126} = (\beta^u/\beta^d) \tan\beta$, $c_T = v_T/(\beta^d v \cos\beta)$ and $c_R = v_R/(\beta^d v \cos\beta)$, respectively. These are the mass matrix relations required by the minimal SO(10) model. In the following in Part I, we set $c_T = 0$ as the first approximation. Except for c_R , which is used to determine the overall neutrino mass scale, this system has fourteen free parameters in total [11], and the strong predictability to the fermion mass matrices. The reasonable results we found are listed in Table 1.

As mentioned above, our resultant neutrino oscillation parameters are sensitive to all the input parameters. In other words, if we use the neutrino oscillation data as the input parameters, the other input, for example, the CP-phase in the CKM matrix can be regarded as the prediction of our model. It is a very interesting observation that the CP-phases listed above are in the region consistent with experiments. The CP-violation in the lepton sector is characterized by the Jarlskog parameter defined as

$$J_{CP} = \text{Im} [U_{e2} U_{\mu 2}^* U_{e3}^* U_{\mu 3}] , \tag{8}$$

where U_{fi} is the MNS matrix element.

Lepton Flavour Violation. It is well known that the SO(10) GUT model possesses a simple mechanism of baryogenesis through the out-of-equilibrium decay of the right-handed neutrinos, namely, the leptogenesis [13]. The lepton asymmetry in the universe is generated by CP-violating out-of-equilibrium decay of the heavy neutrinos, $N \rightarrow \ell_L H_u^*$ and $N \rightarrow \bar{\ell}_L H_u$. The leading contribution is given by the interference between the tree level and one-loop




FIGURE 1. Three mixing angles in the MNS matrix as functions of σ [rad]. The graphs with the highest, middle and lowest peaks correspond to $\sin^2 2\theta_{23}$, $\sin^2 2\theta_{12}$ and $\sin^2 2\theta_{13}$, respectively. The plots of $\sin^2 2\theta_{23}$ and $\sin^2 2\theta_{13}$ have the sharp peaks at $\sigma \sim 3.2$ [rad], while $\sin^2 2\theta_{12}$ has the sharp peak at $\sigma \sim 3.3$ [rad] cited from [12]

TABLE 2. The input values of $\tan\beta$ and the outputs for the CP-violating observables

$\tan\beta$	$\langle m_\nu \rangle_{ee}$ (eV)	J_{CP}	ε
40	0.00122	0.00110	7.39×10^{-5}
45	0.00118	-0.00429	6.80×10^{-5}
50	0.00119	-0.00631	6.50×10^{-5}
55	0.00117	-0.00612	11.2×10^{-5}

level decay amplitudes, and the CP-violating parameter is found to be

$$\varepsilon = \frac{1}{8\pi(Y_\nu Y_\nu^\dagger)_{11}} \sum_{j=2,3} \text{Im} [(Y_\nu Y_\nu^\dagger)_{1j}^2] \{ f(M_{Rj}^2/M_{R1}^2) + 2g(M_{Rj}^2/M_{R1}^2) \}. \quad (9)$$

Here $f(x)$ and $g(x)$ correspond to the vertex and the wave function corrections,

$$\begin{aligned} f(x) &\equiv \sqrt{x} \left[1 - (1+x) \ln \left(\frac{1+x}{x} \right) \right], \\ g(x) &\equiv \frac{\sqrt{x}}{2(1-x)}, \end{aligned} \quad (10)$$

respectively, and both are reduced to $\sim -\frac{1}{2\sqrt{x}}$ for $x \gg 1$. So in this approximation, ε becomes

$$\varepsilon = -\frac{3}{16\pi(Y_\nu Y_\nu^\dagger)_{11}} \sum_{j=2,3} \text{Im} [(Y_\nu Y_\nu^\dagger)_{1j}^2] \frac{M_{R1}}{M_{Rj}}. \quad (11)$$

These quantities are evaluated by using the results presented in Table 1, and the results are listed in Table 2.

Now we turn to the discussion about the rate of the LFV processes and the muon $g-2$. The evidence of the neutrino flavor mixing implies that the lepton flavor of each generation is not individually conserved. Therefore the lepton flavor violating (LFV) processes in the charged-lepton sector such as $\mu \rightarrow e\gamma$, $\tau \rightarrow \mu\gamma$ are allowed. In simply extended models so as to incorporate massive neutrinos into the standard model, the rate of the LFV processes is accompanied

by a highly suppression factor, the ratio of neutrino mass to the weak boson mass, because of the GIM mechanism, and is far out of the reach of the experimental detection. However, in supersymmetric models, the situation is quite different. In this case, soft SUSY breaking parameters can be new LFV sources, and the rate of the LFV processes are suppressed by only the scale of the soft SUSY breaking parameters which is assumed to be the electroweak scale. Thus the huge enhancement occurs compared to the previous case. In fact, the LFV processes can be one of the most important processes as the low-energy SUSY search. we evaluate the rate of the LFV processes in the minimal SUSY SO(10) model, where the neutrino Dirac Yukawa couplings are the primary LFV sources. Although in Ref. [12] various cases with given $\tan\beta = 40 - 55$ have been analyzed, we consider only the case $\tan\beta = 45$ in the following. Our final result is almost insensitive to $\tan\beta$ values in the above range. The predictions of the minimal SUSY SO(10) model necessary for the LFV processes are as follows [12]: with $\sigma = 3.198$ fixed, the right-handed Majorana neutrino mass eigenvalues are found to be (in GeV) $M_{R_1} = 1.64 \times 10^{11}$, $M_{R_2} = 2.50 \times 10^{12}$ and $M_{R_3} = 8.22 \times 10^{12}$, where c_R is fixed so that $\Delta m_{\oplus}^2 = 2 \times 10^{-3} \text{eV}^2$. In the basis where both of the charged-lepton and right-handed Majorana neutrino mass matrices are diagonal with real and positive eigenvalues, the neutrino Dirac Yukawa coupling matrix at the GUT scale is found to be ⁴

$$Y_\nu = \begin{pmatrix} -0.000135 - 0.00273i & 0.00113 + 0.0136i & 0.0339 + 0.0580i \\ 0.00759 + 0.0119i & -0.0270 - 0.00419i & -0.272 - 0.175i \\ -0.0280 + 0.00397i & 0.0635 - 0.0119i & 0.491 - 0.526i \end{pmatrix}. \quad (12)$$

LFV effect most directly emerges in the left-handed slepton mass matrix through the RGEs such as [15]

$$\begin{aligned} \mu \frac{d}{d\mu} (m_{\tilde{\ell}}^2)_{ij} &= \mu \frac{d}{d\mu} (m_{\tilde{\ell}}^2)_{ij} \Big|_{\text{MSSM}} \\ &+ \frac{1}{16\pi^2} (m_{\tilde{\ell}}^2 Y_\nu^\dagger Y_\nu + Y_\nu^\dagger Y_\nu m_{\tilde{\ell}}^2 + 2Y_\nu^\dagger m_{\tilde{\nu}}^2 Y_\nu + 2m_{H_u}^2 Y_\nu^\dagger Y_\nu + 2A_\nu^\dagger A_\nu)_{ij}, \end{aligned} \quad (13)$$

where the first term in the right hand side denotes the normal MSSM term with no LFV. We have found Y_ν explicitly and we can calculate LFV and related phenomena unambiguously [14] In the leading-logarithmic approximation, the off-diagonal components ($i \neq j$) of the left-handed slepton mass matrix are estimated as

$$(\Delta m_{\tilde{\ell}}^2)_{ij} \sim -\frac{3m_0^2 + A_0^2}{8\pi^2} (Y_\nu^\dagger L Y_\nu)_{ij}, \quad (14)$$

where the distinct thresholds of the right-handed Majorana neutrinos are taken into account by the matrix $L = \log[M_{\text{GUT}}/M_{R_i}] \delta_{ij}$.

The effective Lagrangian relevant for the LFV processes ($\ell_i \rightarrow \ell_j \gamma$) and the muon $g-2$ is described as

$$\mathcal{L}_{\text{eff}} = -\frac{e}{2} m_{\ell_i} \bar{\ell}_j \sigma_{\mu\nu} F^{\mu\nu} (A_L^{ji} P_L + A_R^{ji} P_R) \ell_i, \quad (15)$$

where $P_{R,L} = (1 \pm \gamma_5)/2$ is the chirality projection operator, and $A_{L,R}$ are the photon-penguin couplings of 1-loop diagrams in which chargino-sneutrino and neutralino-charged slepton are running. The explicit formulas of $A_{L,R}$ etc. used in our analysis are summarized in [16] [17]. The rate of the LFV decay of charged-leptons is given by

$$\Gamma(\ell_i \rightarrow \ell_j \gamma) = \frac{e^2}{16\pi} m_{\ell_i}^5 \left(|A_L^{ji}|^2 + |A_R^{ji}|^2 \right), \quad (16)$$

while the real diagonal components of $A_{L,R}$ contribute to the anomalous magnetic moments of the charged-leptons such as

$$\delta a_{\ell_i}^{\text{SUSY}} = \frac{g_{\ell_i} - 2}{2} = -m_{\ell_i}^2 \text{Re} [A_L^{ii} + A_R^{ii}]. \quad (17)$$

⁴ We are now reconsidering data fitting with the up to date experimental data and new RGE results. It gives the different values from (12) but the LFV results are not essentially changed.

In order to clarify the parameter dependence of the decay amplitude, we give here an approximate formula of the LFV decay rate [16],

$$\Gamma(\ell_i \rightarrow \ell_j \gamma) \sim \frac{e^2}{16\pi} m_{\ell_i}^5 \times \frac{\alpha_2}{16\pi^2} \frac{\left| \left(\Delta m_{\tilde{\ell}}^2 \right)_{ij} \right|^2}{M_S^8} \tan^2 \beta, \quad (18)$$

where M_S is the average slepton mass at the electroweak scale, and $\left(\Delta m_{\tilde{\ell}}^2 \right)_{ij}$ is the slepton mass estimated in Eq. (14). We can see that the neutrino Dirac Yukawa coupling matrix plays the crucial role in calculations of the LFV processes. We use the neutrino Dirac Yukawa coupling matrix of Eq. (12) in our numerical calculations.

The recent Wilkinson Microwave Anisotropy Probe (WMAP) satellite data [18] provide estimations of various cosmological parameters with greater accuracy. The current density of the universe is composed of about 73% of dark energy and 27% of matter. Most of the matter density is in the form of the CDM, and its density is estimated to be (in 2σ range)

$$\Omega_{\text{CDM}} h^2 = 0.1126_{-0.0181}^{+0.0161}. \quad (19)$$

The parameter space of the CMSSM which allows the neutralino relic density suitable for the cold dark matter has been recently re-examined in the light of the WMAP data [19]. It has been shown that the resultant parameter space is dramatically reduced into the narrow stripe due to the great accuracy of the WMAP data. It is interesting to combine this result with our analysis of the LFV processes and the muon $g-2$. In the case relevant for our analysis, $\tan \beta = 45$, $\mu > 0$ and $A_0 = 0$, we can read off the approximate relation between m_0 and $M_{1/2}$ such as (see Figure 1 in the second paper of Ref. [19])

$$m_0(\text{GeV}) = \frac{9}{28} M_{1/2}(\text{GeV}) + 150(\text{GeV}), \quad (20)$$

along which the neutralino CDM is realized. $M_{1/2}$ parameter space is constrained within the range $300\text{GeV} \leq M_{1/2} \leq 1000\text{GeV}$ due to the experimental bound on the SUSY contribution to the $b \rightarrow s\gamma$ branching ratio and the unwanted stau LSP parameter region. We show $\text{Br}(\mu \rightarrow e\gamma)$ and the muon $g-2$ as functions of $M_{1/2}$ in Fig. ?? and ??, respectively, along the neutralino CDM condition of Eq. (20). We find the parameter region, $560\text{GeV} \leq M_{1/2} \leq 800\text{GeV}$, being consistent with all the experimental data.




FIGURE 2. The branching ratio, a: $\text{Log}_{10}[\text{Br}(\mu \rightarrow e\gamma)]$, b: the SUSY contribution to the muon $g-2$ in units of 10^{-10} , $\delta a_{\mu}^{\text{SUSY}} = \frac{g_{\mu}^2 - 2}{2}$, and c: the electron EDM, $\text{Log}_{10}[|d_e|[\text{e cm}]]$. All these figures are plotted as a function of $M_{1/2}$ (GeV) along the cosmological constraint of Eq. (20).

The semileptonic flavor violation processes were also considered in [20], for instance, for $\tau^- \rightarrow e^-(\mu^-)\pi^0$, $\tau^- \rightarrow e^-(\mu^-)\eta$, $\tau^- \rightarrow e^-(\mu^-)\eta'$, $\tau^- \rightarrow e^-(\mu^-)\rho^0$, $\tau^- \rightarrow e^-(\mu^-)\phi$, $\tau^- \rightarrow e^-(\mu^-)\omega$, etc.

When the KamLAND data [21] was released, the results in [12] were found to be deviated by 3σ from the observations. Afterward this minimal SO(10) was modified by many authors, using the so-called type-II see-saw mechanism [22] and/or considering a **120** Higgs coupling to the matter in addition to the **126** Higgs [23]. Based on an elaborate input data scan [25], [24], it has been shown that the minimal SO(10) is essentially consistent with low energy data of fermion masses and mixing angles. The importance of the threshold corrections was also discussed in [26]

Higgs Superpotential. On the other hand, it has been long expected to construct a concrete Higgs sector of the minimal $SO(10)$ model. The simplest Higgs superpotential at the renormalizable level is given by [28], [29], [30]

$$W = m_1 \Phi^2 + m_2 \Delta \bar{\Delta} + m_3 H^2 + \lambda_1 \Phi^3 + \lambda_2 \Phi \Delta \bar{\Delta} + \lambda_3 \Phi \Delta H + \lambda_4 \Phi \bar{\Delta} H, \quad (21)$$

where $\Phi = \mathbf{210}$, $\Delta = \mathbf{126}$, $\bar{\Delta} = \overline{\mathbf{126}}$ and $H = \mathbf{10}$. The interactions of $\mathbf{210}$, $\overline{\mathbf{126}}$, $\mathbf{126}$ and $\mathbf{10}$ lead to some complexities in decomposing the GUT representations to the MSSM and in getting the low energy mass spectra. Particularly, the CG coefficients corresponding to the decompositions of $SO(10) \rightarrow SU(3)_C \times SU(2)_L \times U(1)_Y$ have to be found. This problem was first attacked by X. G. He and S. Meljanac [31] and further by D. G. Lee [29] and by J. Sato [32]. But they did not present the explicit form of mass matrices for a variety of Higgs fields and also did not perform a formulation of the proton life time analysis. This is very labourious work and it is indispensable for the data fit of low energy physics. We completed that program in [33] (See also [34], [35]). This construction gives some constraints among the vacuum expectation values (VEVs) of several Higgs multiplets, which give rise to a trouble in the gauge coupling unification [24]. The trouble comes from the fact that the observed neutrino oscillation data suggests the right-handed neutrino mass around 10^{13-14} GeV, which is far below the GUT scale. Indeed (21) contains five directions which are singlets under $SU(3)_C \times SU(2)_L \times U(1)_Y$. Three of them are included in $\mathbf{210}$,

$$\hat{\phi}_1 = (1234), \quad (22)$$

$$\hat{\phi}_2 = (5678 + 5690 + 7890), \quad (23)$$

$$\hat{\phi}_3 = (12 + 34)(56 + 78 + 90). \quad (24)$$

one in $\mathbf{126}$ and

$$\hat{v}_R = (13579), \quad (25)$$

and one in $\overline{\mathbf{126}}$

$$\hat{\bar{v}}_R = (24680). \quad (26)$$

Due to the D-flatness condition the VEVs v_R and $\bar{v}_R$ are equal (21),

$$v_R = \bar{v}_R. \quad (27)$$

This intermediate scale is provided by Higgs field VEV, and several Higgs multiplets are expected to have their masses around the intermediate scale and contribute to the running of the gauge couplings.

We write down the VEV conditions which preserve supersymmetry, with respect to the directions $\hat{\phi}_1$, $\hat{\phi}_2$, $\hat{\phi}_3$, and $\hat{v}_R$, respectively.

$$2m_1 \phi_1 + 3\lambda_1 \frac{\phi_3^2}{6\sqrt{6}} + \lambda_2 \frac{v_R^2}{10\sqrt{6}} = 0, \quad (28)$$

$$2m_1 \phi_2 + 3\lambda_1 \left(\frac{\phi_2^2 + \phi_3^2}{9\sqrt{2}} \right) + \lambda_2 \frac{v_R^2}{10\sqrt{2}} = 0, \quad (29)$$

$$2m_1 \phi_3 + 3\lambda_1 \left(\frac{\phi_1 \phi_3}{3\sqrt{6}} + \frac{\sqrt{2} \phi_2 \phi_3}{9} \right) + \lambda_2 \frac{v_R^2}{10} = 0, \quad (30)$$

$$m_2 + \lambda_2 \left(\frac{\phi_1}{10\sqrt{6}} + \frac{\phi_2}{10\sqrt{2}} + \frac{\phi_3}{10} \right) = 0. \quad (31)$$

Eliminating v_R^2 , ϕ_1 and ϕ_3 from Eqs. (28)–(31), one obtains a fourth-order equation in ϕ_2 . The corresponding fourth-order polynomial in ϕ_2 factorizes into a linear and a cubic term in ϕ_2 . Linear term gives the solution of the fourth-order equation which is very simple, $\phi_2 = -3\sqrt{2}m_2/\lambda_2$, but it preserves the $SU(5)$ symmetry. Therefore, it is physically not interesting. The cubic term solutions lead to the true $SU(3)_C \times SU(2)_L \times U(1)_Y$ symmetry. Here we consider only the solutions with $|v_R| \neq 0$. Eliminating $v_R \cdot \bar{v}_R$, ϕ_1 and ϕ_2 from Eqs. (28)–(31), one obtains a fourth-order equation in ϕ_3 ,

$$\left(\phi_3 + \frac{\mathcal{M}_2}{10} \right) \left\{ 8\phi_3^3 - 15\mathcal{M}_1\phi_3^2 + 14\mathcal{M}_1^2\phi_3 - 3\mathcal{M}_1^3 + (\phi_3 - \mathcal{M}_1)^2 \mathcal{M}_2 \right\} = 0, \quad (32)$$

where

$$\mathcal{M}_1 \equiv 12 \left(\frac{m_1}{\lambda_1} \right), \mathcal{M}_2 \equiv 60 \left(\frac{m_2}{\lambda_2} \right). \quad (33)$$

Any solution of the cubic equation in ϕ_3 is accompanied by the solutions

$$\phi_1 = -\frac{\phi_3 (\mathcal{M}_1^2 - 5\phi_3^2)}{\sqrt{6} (\mathcal{M}_1 - \phi_3)^2}, \quad (34)$$

$$\phi_2 = -\frac{1}{\sqrt{2}} \frac{(\mathcal{M}_1^2 - 2\mathcal{M}_1\phi_3 - \phi_3^2)}{(\mathcal{M}_1 - \phi_3)}, \quad (35)$$

$$v_R \cdot \overline{v_R} = \frac{5}{3} \left(\frac{\lambda_1}{\lambda_2} \right) \frac{\phi_3 (\mathcal{M}_1 - 3\phi_3) (\mathcal{M}_1^2 + \phi_3^2)}{(\mathcal{M}_1 - \phi_3)^2}. \quad (36)$$

The linear term gives the solution of the fourth-order equation (32) which is very simple, $\phi_3 = -6 \left(\frac{m_2}{\lambda_2} \right)$. It leads to $\phi_1 = -\sqrt{6} \left(\frac{m_2}{\lambda_2} \right)$, $\phi_2 = -3\sqrt{2} \left(\frac{m_2}{\lambda_2} \right)$ and $\sqrt{v_R \cdot \overline{v_R}} = \sqrt{60} \left(\frac{m_2}{\lambda_2} \right) \sqrt{2 \left(\frac{m_1}{m_2} \right) - 3 \left(\frac{\lambda_1}{\lambda_2} \right)}$. This solution preserves the $SU(5)$ symmetry. Therefore, it is physically not interesting. Then we proceed to the most important part of the $SO(10)$ GUT. We can not show the detail of the scenario but only show the essential part of it [33].

Would-be Nambu-Goldstone bosons. At first, we list the quantum numbers of the would-be NG modes under $SU(3)_C \times SU(2)_L \times U(1)_Y$.

- $[(\overline{\mathbf{3}}, \mathbf{2}, \frac{5}{6}) \oplus (\mathbf{3}, \mathbf{2}, -\frac{5}{6})]$,
- $[(\overline{\mathbf{3}}, \mathbf{2}, -\frac{1}{6}) \oplus (\mathbf{3}, \mathbf{2}, \frac{1}{6})]$,
- $[(\overline{\mathbf{3}}, \mathbf{1}, -\frac{2}{3}) \oplus (\mathbf{3}, \mathbf{1}, \frac{2}{3})]$,
- $[(\mathbf{1}, \mathbf{1}, 1) \oplus (\mathbf{1}, \mathbf{1}, -1)]$,
- $[(\mathbf{1}, \mathbf{1}, 0)]$.

Total number of the NG degrees of freedom is : $\mathbf{12} + \mathbf{12} + \mathbf{6} + \mathbf{2} + \mathbf{1} = \mathbf{33}$. The cubic term solutions lead to the true $SU(3)_C \times SU(2)_L \times U(1)_Y$ symmetry. (36) gives heavy right-handed neutrino, and the coefficient of (7) is also written in terms of ϕ_3 .

Electroweak Higgs doublet. In the standard picture of the electroweak symmetry breaking, we have the Higgs doublets which give masses to the matter. These masses should be less than or equal to the electroweak scale. Since we approximate the electroweak scale as zero, we must impose a constraint that the mass matrix should have one zero eigenvalue.

We define

$$H_u^{10} \equiv H_{(1,2,2)}^{(1,2,\frac{1}{2})}, \overline{\Delta}_u \equiv \overline{\Delta}_{(15,2,2)}^{(1,2,\frac{1}{2})}, \Delta_u \equiv \Delta_{(15,2,2)}^{(1,2,\frac{1}{2})}, \Phi_u \equiv \Phi_{(10,2,2)}^{(1,2,\frac{1}{2})}. \quad (37)$$

and

$$H_d^{10} \equiv H_{(1,2,2)}^{(1,2,-\frac{1}{2})}, \overline{\Delta}_d \equiv \overline{\Delta}_{(15,2,2)}^{(1,2,-\frac{1}{2})}, \Delta_d \equiv \Delta_{(15,2,2)}^{(1,2,-\frac{1}{2})}, \Phi_d \equiv \Phi_{(10,2,2)}^{(1,2,-\frac{1}{2})}. \quad (38)$$

In the basis $\{H_u^{10}, \overline{\Delta}_u, \Delta_u, \Phi_u\}$, the mass matrix is written as

$$M_{\text{doublet}} \equiv \begin{pmatrix} 2m_3 & \frac{\lambda_3\phi_1}{\sqrt{10}} - \frac{\lambda_3\phi_3}{2\sqrt{5}} & -\frac{\lambda_4\phi_2}{\sqrt{10}} - \frac{\lambda_4\phi_3}{2\sqrt{5}} & \frac{\lambda_4\overline{v_R}}{\sqrt{5}} \\ \frac{\lambda_4\phi_2}{\sqrt{10}} - \frac{\lambda_4\phi_3}{2\sqrt{5}} & m_2 + \frac{\lambda_2\phi_2}{15\sqrt{2}} - \frac{\lambda_2\phi_3}{30} & 0 & 0 \\ -\frac{\lambda_3\phi_2}{\sqrt{10}} - \frac{\lambda_3\phi_3}{2\sqrt{5}} & 0 & m_2 + \frac{\lambda_2\phi_2}{15\sqrt{2}} + \frac{\lambda_2\phi_3}{30} & -\frac{\lambda_2\overline{v_R}}{10} \\ \frac{\lambda_3\overline{v_R}}{\sqrt{5}} & 0 & -\frac{\lambda_2\overline{v_R}}{10} & 2m_1 + \frac{\lambda_1\phi_2}{\sqrt{2}} + \frac{\lambda_1\phi_3}{2} \end{pmatrix}. \quad (39)$$

The corresponding mass terms of the superpotential read

$$W_m = (H_u^{10}, \overline{\Delta}_u, \Delta_u, \Phi_u) M_{\text{doublet}} (H_d^{10}, \Delta_d, \overline{\Delta}_d, \Phi_d)^T. \quad (40)$$

The requirement of the existence of a zero mode leads to the following condition.

$$\det M_{\text{doublet}} = 0. \quad (41)$$

For instance, in case of $\lambda_3 = 0$, $m_2 + \frac{\lambda_2 \phi_2}{15\sqrt{2}} - \frac{\lambda_2 \phi_3}{30} = 0$, we obtain a special solution to Eq. (41), while it keeps a desirable vacuum and it does not produce any additional massless fields. However, we proceed our arguments hereafter without using this special solution.

We can diagonalize the mass matrix, M_{doublet} by a bi-unitary transformation.

$$U^* M_{\text{doublet}} V^\dagger = \text{diag}(0, M_1, M_2, M_3). \quad (42)$$

Then the mass eigenstates are written as

$$\begin{aligned} (H_u, h_u^1, h_u^2, h_u^3) &= (H_u^{10}, \bar{\Delta}_u, \Delta_u, \Phi_u) U^\top, \\ (H_d, h_d^1, h_d^2, h_d^3) &= (H_d^{10}, \Delta_d, \bar{\Delta}_d, \Phi_d) V^\top. \end{aligned} \quad (43)$$

Here H_u, H_d are MSSM light Higgs doublets. We get the explicit form of U and V from (39), and thus we can connect the oacillation data with the GUT Yukawa coupling. Thus the intermediate energy scales are severely constrained from the low energy neutrino data, and the gauge coupling unification at the GUT scale may be spoiled.

This fact has been explicitly shown in [36], where the gauge couplings are not unified any more and even the $SU(2)$ gauge coupling blows up below the GUT scale. Thus the detail analyses of superpotential was the great progress but it reveals the unambiguous detail of structure, which reveals also pathologies.

However, this is easily remedied by the addition of **120** Higgs in Yukawa coupling [37]. We mean that the dominant part may be governed by the minimal $SO(10)$ but such generalization does not spoil the renormalizable $SO(10)$ GUT yet.

General Higgs superpotential. Also we may consider the more general superpotential for completeness [27].

$$\begin{aligned} W &= \frac{1}{2} m_1 \Phi^2 + m_2 \bar{\Delta} \Delta + \frac{1}{2} m_3 H^2 \\ &+ \frac{1}{2} m_4 A^2 + \frac{1}{2} m_5 E^2 + \frac{1}{2} m_6 D^2 \\ &+ \lambda_1 \Phi^3 + \lambda_2 \Phi \bar{\Delta} \Delta + (\lambda_3 \Delta + \lambda_4 \bar{\Delta}) H \Phi \\ &+ \lambda_5 A^2 \Phi - i \lambda_6 A \bar{\Delta} \Delta + \frac{\lambda_7}{120} \epsilon A \Phi^2 \\ &+ E \left(\lambda_8 E^2 + \lambda_9 A^2 + \lambda_{10} \Phi^2 + \lambda_{11} \Delta^2 + \lambda_{12} \bar{\Delta}^2 + \lambda_{13} H^2 \right) \\ &+ D^2 (\lambda_{14} E + \lambda_{15} \Phi) \\ &+ D \{ \lambda_{16} H A + \lambda_{17} H \Phi + (\lambda_{18} \Delta + \lambda_{19} \bar{\Delta}) A + (\lambda_{20} \Delta + \lambda_{21} \bar{\Delta}) \Phi \}, \end{aligned} \quad (44)$$

Here $A = \mathbf{45}$, $\Delta = \mathbf{126}$, $\Phi = \mathbf{210}$ and $E = \mathbf{54}$ irreps. For general coupling constants $\lambda_1, \dots, \lambda_{21}$, $m_1, \dots, m_8$, the solutions with higher symmetries are specified by following relations. Solutions with higher symmetries are characterized by:

1. $SU(5) \times U(1)_X$ and $(SU(5) \times U(1))^{\text{flipped}}$ symmetry solutions

$$\begin{cases} E &= v_R = 0, \\ \Phi_1 &= \frac{\epsilon}{\sqrt{6}} \Phi_3, \quad \Phi_2 = \frac{\epsilon}{\sqrt{2}} \Phi_3, A_1 = \frac{2\epsilon}{\sqrt{6}} A_2, \end{cases} \quad (45)$$

where $\epsilon = 1$ and $\epsilon = -1$ correspond to the $SU(5) \times U(1)_X$ symmetric vacua and $(SU(5) \times U(1))^{\text{flipped}}$ symmetric vacua, respectively.

2. $SU(5)$ symmetry solutions

$$\begin{cases} E &= 0, \\ \Phi_1 &= \frac{1}{\sqrt{6}} \Phi_3, \quad \Phi_2 = \frac{1}{\sqrt{2}} \Phi_3, \quad A_1 = \frac{2}{\sqrt{6}} A_2, \quad v_R \neq 0. \end{cases} \quad (46)$$

3. $G_{422} \equiv SU(4) \times SU(2)_L \times SU(2)_R$ symmetry solutions

$$\begin{cases} \Phi_2 &= \Phi_3 = A_1 = A_2 = v_R = 0, \\ \Phi_1 &\neq 0, \quad E \neq 0. \end{cases} \quad (47)$$

4. $G_{3221} \equiv SU(3)_C \times SU(2)_L \times SU(2)_R \times U(1)_{B-L}$ symmetry solutions

$$\begin{cases} \Phi_3 = A_1 = v_R = 0, \\ \Phi_1 \neq 0, \quad \Phi_2 \neq 0, \quad A_2 \neq 0, \quad E \neq 0. \end{cases} \quad (48)$$

5. $G_{421} \equiv SU(4) \times SU(2)_L \times U(1)$ symmetry solutions

$$\begin{cases} \Phi_2 = \Phi_3 = A_2 = v_R = 0 \\ \Phi_1 \neq 0, \quad A_1 \neq 0, \quad E \neq 0. \end{cases} \quad (49)$$

6. $G_{3211} \equiv SU(3)_C \times SU(2)_L \times U(1)_R \times U(1)_{B-L}$ symmetry solutions

$$\begin{cases} v_R = 0, \\ \Phi_i \neq 0 \quad (i = 1, 2, 3), \quad A_i \neq 0 \quad (i = 1, 2), \quad E \neq 0. \end{cases} \quad (50)$$

The higher symmetry solutions given in Eqs. (45)-(50) lead to the crucial consistency checks for all results in this paper. In this talk, however, we need the alternative approaches.

In order to avoid this trouble more drastically and keep the successful gauge coupling unification as usual, it is desirable that all Higgs multiplets have masses around the GUT scale, but some Higgs fields develop VEVs at the intermediate scale. More Higgs multiplets and some parameter tuning in the Higgs sector are necessary to realize such a situation.

In addition to the issue of the gauge coupling unification, the minimal SO(10) model potentially suffers from the problem that the gauge coupling blows up around the GUT scale. This is because the model includes many Higgs multiplets of higher dimensional representations.

According to the line of thoughts from (1) to (2), it was natural to consider

$$L_{GUT} = L''_{ren} + \frac{L_3}{\Lambda_3}. \quad (51)$$

up to M_{PL} . Here $\Lambda_3 = O(M_{Pl})$ and gravitation (spacetime structure) appears as a subdominant term. However the blow-up before M_{PL} problem shows that such scheme does not exist in its naive sense.

The minimal SO(10) model also is faced on the fast proton decay [39]

These facts strongly (but not indispensably) suggest the presence of extra dimensions, which gives not only solve the above problems but also new insights for SUSY breaking mechanism [40]

PART II—SO(10) GUT IN 5D

In this Part we propose a solution to the problem of the minimal SO(10) discussed in Part I.

Minimal SO(10) model in a warped extra dimension

We consider the minimal SUSY SO(10) model in the following 5D warped geometry⁵

$$ds^2 = e^{-2kr_c|y|} \eta_{\mu\nu} dx^\mu dx^\nu - r_c^2 dy^2, \quad (52)$$

for $-\pi \leq y \leq \pi$, where k is the AdS curvature, and r_c and y are the radius and the angle of S^1 , respectively [41]. The most important feature of the warped extra dimension model is that the mass scale of the IR brane is warped down to a low scale by the warp factor [41], $\omega \equiv e^{-kr_c\pi}$, in four dimensional effective theory. For simplicity, we take the cutoff of the original five dimensional theory and the AdS curvature as $M_5 \simeq k \simeq M_{PL}$, the four dimensional Planck mass, and so we obtain the effective cutoff scale as $\Lambda_{IR} = \omega M_{PL}$ in effective four dimensional theory. Now let us take the warp

⁵ This part is based on the work, “Solving problems of 4D minimal SO(10) model in a warped extra dimension”, T. Fukuyama, T. Kikuchi and N. Okada, Phys.Rev.**D75** 075020 (2007) [Archive: hep-ph/0702048].

factor so as for the GUT scale to be the effective cutoff scale $M_{\text{GUT}} = \Lambda_{\text{IR}} = \omega M_{\text{PL}}$ [42]. As a result, we can realize, as four dimensional effective theory, the minimal SUSY SO(10) model with the effective cutoff at the GUT scale.

Before going to a concrete setup of the minimal SO(10) model in the warped extra dimension, let us see Lagrangian for the hypermultiplet in the bulk,

$$\begin{aligned} \mathcal{L} = & \int dy \left\{ \int d^4\theta r_c e^{-2kr_c|y|} (H^\dagger e^{-V} H + H^c e^V H^{c\dagger}) \right. \\ & \left. + \int d^2\theta e^{-3kr_c|y|} H^c \left[\partial_y - \left(\frac{3}{2} - c \right) kr_c \varepsilon(y) - \frac{\chi}{\sqrt{2}} \right] H + h.c. \right\}, \end{aligned} \quad (53)$$

where c is a dimensionless parameter, $\varepsilon(y) = y/|y|$ is the step function, H , H^c is the hypermultiplet charged under some gauge group, and V , χ are the vector multiplet and the adjoint chiral multiplets, which form an $N = 2$ SUSY gauge multiplet. Z_2 parity for H and V is assigned as even, while odd for H^c and χ .

When the gauge symmetry is broken down, it is generally possible that the adjoint chiral multiplet develops its VEV [46]. Since its Z_2 parity is odd, the VEV has to take the form,

$$\langle \Sigma \rangle = 2\alpha kr_c \varepsilon(y), \quad (54)$$

where the VEV has been parameterized by a parameter α . In this case, the zero mode wave function of H satisfies the following equation of motion:

$$\left[\partial_y - \left(\frac{3}{2} - c + \alpha \right) kr_c \varepsilon(y) \right] H = 0 \quad (55)$$

which yields

$$H = \frac{1}{\sqrt{N}} e^{(3/2 - c + \alpha)kr_c|y|} h(x^\mu), \quad (56)$$

where $h(x^\mu)$ is the chiral multiplet in four dimensions.

Lagrangian for a chiral multiplets on the IR brane is given by

$$\mathcal{L}_{\text{IR}} = \int d^4\theta \omega^\dagger \omega \Phi^\dagger \Phi + \left[\int d^2\theta \omega^3 W(\Phi) + h.c. \right], \quad (57)$$

where we have omitted the gauge interaction part for simplicity. If it is allowed by the gauge invariance, we can write the interaction term between fields in the bulk and on the IR brane,

$$\mathcal{L}_{\text{int}} = \int d^2\theta \omega^3 \frac{Y}{\sqrt{M_5}} \Phi^2 H(y = \pi) + h.c., \quad (58)$$

where Y is a Yukawa coupling constant, and M_5 is the five dimensional Planck mass (we take $M_5 \sim M_{\text{PL}}$ as mentioned above, for simplicity). Rescaling the brane field $\Phi \rightarrow \Phi/\omega$ to get the canonically normalized kinetic term and substituting the zero-mode wave function of the bulk fields, we obtain Yukawa coupling constant in effective four dimensional theory as

$$Y_{4D} \sim Y \quad (59)$$

if $e^{(1/2 - c + \alpha)kr_c\pi} \gg 1$, while

$$Y_{4D} \sim Y \times e^{(1/2 - c + \alpha)kr_c\pi} \ll Y, \quad (60)$$

for $e^{(1/2 - c + \alpha)kr_c\pi} \ll 1$. In the latter case, we obtain a suppression factor since H is localized around the UV brane.

Now we give a simple setup of the minimal SO(10) model in the warped extra dimension. We put all **16** matter multiplets on the IR ($y = \pi$) brane, while the Higgs multiplets **10** and **$\overline{126}$** are assumed to live in the bulk. In Eq. (58), replacing the brane field into the matter multiplets and the bulk field into the Higgs multiplets, we obtain Yukawa couplings in the minimal SO(10) model. The Lagrangian for the bulk Higgs multiplets are given in the same form as Eq. (53), where χ is the SO(10) adjoint chiral multiplet, **45**. As discussed above, since the SO(10) gauge group is

broken down to the SM one, some components in χ which is singlet under the SM gauge group can in general develop VEVs. Here we consider a possibility that the $U(1)_X$ component in the adjoint $\chi = \mathbf{45}$ under the decomposition $SO(10) \supset SU(5) \times U(1)_X$ has a non-zero VEV,

$$\mathbf{45} = \mathbf{1}_0 \oplus \mathbf{10}_{+4} \oplus \overline{\mathbf{10}}_{-4} \oplus \mathbf{24}_0 .$$

The $\overline{\mathbf{126}}$ Higgs multiplet are decomposed under $SU(5) \times U(1)_X$ as

$$\overline{\mathbf{126}} = \mathbf{1}_{+10} \oplus \mathbf{5}_{+2} \oplus \overline{\mathbf{10}}_{+6} \oplus \mathbf{15}_{-6} \oplus \overline{\mathbf{45}}_{-2} \oplus \mathbf{50}_{+2} .$$

In this decomposition, the coupling between a bulk Higgs multiplet and the $U(1)_X$ component in χ is proportional to $U(1)_X$ charge,

$$\mathcal{L}_{int} \supset \frac{1}{2} \int d^2\theta \omega^3 Q_X \langle \Sigma_X \rangle H^c H + h.c. , \quad (61)$$

and thus each component effectively obtains the different bulk mass term,

$$\left(\frac{3}{2} - c \right) k r_c + \frac{1}{2} Q_X \langle \Sigma_X \rangle , \quad (62)$$

where Q_X is the $U(1)_X$ charge of corresponding Higgs multiplet, and Σ_X is the scalar component of the $U(1)_X$ gauge multiplet ($\mathbf{1}_0$). Now we obtain different configurations of the wave functions for these Higgs multiplets. Since the $\mathbf{1}_{+10}$ Higgs has a large $U(1)_X$ charge relative to other Higgs multiplets, we can choose parameters c and $\langle \Sigma_X \rangle$ so that Higgs doublets are mostly localized around the IR brane while the $\mathbf{1}_{+10}$ Higgs is localized around the UV brane. Therefore, we obtain a suppression factor as in Eq. (60) for the effective Yukawa coupling between the Higgs and right-handed neutrinos. In effective four dimensional description, the GUT mass matrix relation is partly broken down, and the last term in Eq. (6) is replaced into

$$Y_{126}^{ij} \nu_R \rightarrow Y_{126}^{ij} (\varepsilon \nu_R) , \quad (63)$$

where ε denotes the suppression factor. By choosing an appropriate parameters so as to give $\varepsilon = 10^{-2} - 10^{-3}$, we can take $\nu_R \sim M_{GUT}$ and keep the successful gauge coupling unification in the MSSM.

Thus, in order to solve the problems in 4D, we have considered the minimal $SO(10)$ model in the warped extra dimension. As a simple setup, we have assumed that matter multiplets reside on the IR brane while the Higgs multiplets reside in the bulk. The warped geometry leads to a low scale effective cutoff in effective four dimensional theory, and we fix it at the GUT scale. Therefore, the four dimensional minimal $SO(10)$ model is realized as the effective theory with the GUT scale cutoff.

However, it gives rise another problem: $SO(10)$ is anomaly free theory and there appears no D term as far as we consider spontaneously broken scenario. So we can not cancel D term caused by the vev of $\mathbf{45}$ either at bulk and branes. We did not propose the mechanism how $\mathbf{45}$ has vev. Also it was not clear about the dangerous proton decay. Of course we overlooked some point and we do not exclude this scenario.

In the next section we consider another possibility, orbifold GUT model. We break the original $N=2$ (in the sense of 4D) SUSY $SO(10)$ invariant theory into $SU(4)_C \times SU(2)_L \times SU(2)_R$ (hereafter PS for short) not spontaneously but by the boundary conditions in the orbifold $S^1/Z_2 \times Z'_2$. If we consider extra dimensions, the chiral fields need orbifold like compactification [53]. There are so many papers in this region. We consider the most simple and clearcut scenario in this paper.

Orbifold GUT

We consider a SUSY $SO(10)$ SUSY GUT in 5D orbifold.⁶ Usually $SO(10)$ is considered in six dimension, whereas $SU(5)$ in 5 dimension. This is because we need at least two projections for $SO(10)$ down to SM [53][54] if we break the symmetry only through boundary conditions. It should be remarked that even in this case we need the Higgs mechanism to break SM to $SU(3)_c \times U(1)_{em}$. So dimensionality 6 does not have definite meaning.

⁶ This part is based on the work by T. Fukuyama and N. Okada, "A simple $SO(10)$ GUT in five dimensions" [arXiv:hep-ph/0803.1758].

It is very important that in the PS brane we can discard $(\mathbf{6}, \mathbf{2}, \mathbf{1})$ as mentioned in the previous section, and harmful proton decay is circumvented without boundary condition. As mentioned, orbifold is essential for chiral dynamics in extra dimension. However, more extra dimensions are not indispensable in GUT framework. Our set up also indicates why the PS brane is visible brane, where the PS (and not $\text{SO}(10)$) is broken by Higgs mechanism.

In 4D language, 5D vector multiplet consists of $N=1$ vector supermultiplet V and an $N=1$ chiral multiplet Ψ . In $\text{SO}(10)$ V is $\mathbf{45}$.

One extra dimension is compactified on the orbifold $S^1/Z_2 \times Z'_2$ [55]. That is, $N=2$ $\text{SO}(10)$ invariant action in 5D is decomposed into $N=1$ $\text{SO}(10)$ invariant $y = 0$ brane and the PS invariant brane at $y = \pi R/2$ by the boundary conditions of bulk gauge (See Table I). We do consider neither matter nor Higgs in the bulk. For the different set up, see [56] [57] [58].

TABLE 3. P and P' assignment and masses ($n \geq 0$) of fields in the vector multiplet (V, Φ) under the PS group. P' even V contains the PS gauge bosons.

(P, P')	field	mass
$(+, +)$	$V(15, 1, 1), V(1, 3, 1), V(1, 1, 3)$	$\frac{2n}{R}$
$(+, -)$	$V(6, 2, 2)$	$\frac{(2n+1)}{R}$
$(-, +)$	$\Phi(6, 2, 2)$	$\frac{(2n+1)}{R}$
$(-, -)$	$\Phi(15, 1, 1), \Phi(1, 3, 1), \Phi(1, 1, 3)$	$\frac{(2n+2)}{R}$

TABLE 4. Particle contents on the PS brane. Here, we impose the left-right symmetry.

	brane at $y = \pi R/2$
Matter Multiplets	$\psi_i = F_{Li} \oplus F_{Ri} \quad (i = 1, 2, 3)$
Higgs Multiplets	$(\mathbf{1}, \mathbf{2}, \mathbf{2})_H, (\mathbf{1}, \mathbf{2}, \mathbf{2})'_H, (\mathbf{15}, \mathbf{1}, \mathbf{1})_H, (\mathbf{6}, \mathbf{1}, \mathbf{1})_H$ $(\mathbf{4}, \mathbf{1}, \mathbf{2})_H, (\bar{\mathbf{4}}, \mathbf{1}, \mathbf{2})_H, (\mathbf{4}, \mathbf{2}, \mathbf{1})_H, (\bar{\mathbf{4}}, \mathbf{2}, \mathbf{1})_H$

The PS invariance in the PS brane is broken by the Higgs mechanism down to the Standard model by the PS subgroup of $(\mathbf{4}, \mathbf{1}, \mathbf{2})_H$ (H_R terms in (70)). In the $\text{SO}(10)$ model[59], the left- (right-) handed fermions $\psi_{L(R)i}$ in a given i -th generation are assigned to a single irreducible $\mathbf{16}$. Since $\mathbf{16} \times \mathbf{16} = \mathbf{10}_S + \mathbf{120}_A + \mathbf{126}_S$, the fermion masses are generated when the Higgs fields of $\mathbf{10}$, and $\mathbf{120}$, and $\mathbf{126}$ dimensional $\text{SO}(10)$ representation (denoted by ϕ_{10} , ϕ_{120} , and ϕ_{126} , respectively) develop nonvanishing expectation values. Their decomposition under $SU(4) \times SU_L(2) \times SU_R(2)$ are given by

$$\begin{aligned}
\mathbf{10} &= (\mathbf{6}, \mathbf{1}, \mathbf{1}) + (\mathbf{1}, \mathbf{2}, \mathbf{2}), \\
\mathbf{120} &= (\mathbf{15}, \mathbf{2}, \mathbf{2}) + (\mathbf{6}, \mathbf{3}, \mathbf{1}) + (\bar{\mathbf{6}}, \mathbf{1}, \mathbf{3}) + (\mathbf{1}, \mathbf{2}, \mathbf{2}) + (\mathbf{20}, \mathbf{1}, \mathbf{1}), \\
\mathbf{126} &= (\mathbf{10}, \mathbf{3}, \mathbf{1}) + (\bar{\mathbf{10}}, \mathbf{1}, \mathbf{3}) + (\mathbf{15}, \mathbf{2}, \mathbf{2}) + (\bar{\mathbf{6}}, \mathbf{1}, \mathbf{1}).
\end{aligned} \tag{64}$$

On the other hand, the fermion field of 16-dimensinal $\text{SO}(10)$ representation is decomposed as

$$\mathbf{16} = (\mathbf{4}, \mathbf{2}, \mathbf{1}) + (\bar{\mathbf{4}}, \mathbf{1}, \mathbf{2}). \tag{65}$$

With respect to $SU(4) \times SU_L(2) \times SU_R(2)$, the left - and right- handed quarks and leptons of a given i-th generation are assigned as

$$\begin{pmatrix} u_r & u_y & u_b & \nu_e \\ d_r & d_y & d_b & e \end{pmatrix}_{L(R)} \equiv F_{L(R)1}, \quad (66)$$

$F_{L(R)2}$ and $F_{L(R)3}$ are likewise defined for the 2nd and 3rd generations. Note that their transformation properties are $F_{Li} = (\mathbf{4}, \mathbf{2}, \mathbf{1})$ and $F_{Ri} = (\mathbf{4}, \mathbf{1}, \mathbf{2})$ and that $(F_{Li} + \overline{F_{Ri}})$ yields the **16** of $SO(10)$. Since $(\mathbf{4}, \mathbf{2}, \mathbf{1}) \times (\overline{\mathbf{4}}, \mathbf{1}, \mathbf{2}) = (\mathbf{15}, \mathbf{2}, \mathbf{2}) + (\mathbf{1}, \mathbf{2}, \mathbf{2})$, the Dirac masses for quarks and leptons are generated by $(\mathbf{15}, \mathbf{2}, \mathbf{2})_H + (\mathbf{1}, \mathbf{2}, \mathbf{2})_H$.

From $\tau - b$ unification at GUT scale the third generation is described by $(\mathbf{1}, \mathbf{2}, \mathbf{2})$ of H_{10} . The deviations of the first and second generations are complimented by the $(\mathbf{15}, \mathbf{2}, \mathbf{2})_H$ (here it is constructed from $(\mathbf{15}, \mathbf{1}, \mathbf{1}) \times (\mathbf{1}, \mathbf{2}, \mathbf{2})'$: see Table 4). and

$$\begin{aligned} (\mathbf{15}, \mathbf{2}, \mathbf{2}) &= (\mathbf{1}, \mathbf{2}, \mathbf{1}/2) + (\mathbf{1}, \mathbf{2}, -\mathbf{1}/2) + (\mathbf{8}, \mathbf{2}, \mathbf{1}/2) + (\mathbf{8}, \mathbf{2}, -\mathbf{1}/2) + (\mathbf{3}, \mathbf{2}, \mathbf{1}/6) + (\overline{\mathbf{3}}, \mathbf{2}, -\mathbf{1}/6) \\ &+ (\mathbf{3}, \mathbf{2}, \mathbf{7}/6) + (\overline{\mathbf{3}}, \mathbf{2}, -\mathbf{7}/6) \end{aligned} \quad (67)$$

It is remarkable that the component $(\mathbf{6}, \mathbf{1}, \mathbf{1})$ which was harmful for $SO(10)$ invariant Yukawa coupling does not appear [60], and therefore dimension five operator too.

The spectrum of the PS phases were fully discussed in [33]. The third, fourth octet and the last two triplets in (67) becomes massive by the coupling with the counter parts of another $(\mathbf{15}, \mathbf{2}, \mathbf{2})$. The fifth and sixth triplet become NG bosons.

$SU(4)$ adjoint 15 have a basis, $\text{diag}(1, 1, 1, -3)$ so as to satisfy the traceless condition. Putting leptons into the 4th color, we get, so called, A ϵ Georgi-JarlskogA ϵ factor, -3 for leptons. Unlike the case of $SO(10)$, the mass matrix forms of M_L and M_R belong to groups different to each other and different from charged fermions. We may write it in the more familiar forms

$$\begin{aligned} M_u &= c_{10}M_{1,2,2} + c_{15}M_{15,2,2}, \\ M_d &= M_{1,2,2} + M_{15,2,2}, \\ M_D &= c_{10}M_{1,2,2} - 3c_{15}M_{15,2,2}, \\ M_e &= M_{1,2,2} - 3M_{15,2,2}, \\ M_L &= c_L M_{10,3,1}, \\ M_R &= c_R M_{10,1,3}, \end{aligned} \quad (68)$$

Here the effective $(\mathbf{10}, \mathbf{1}, \mathbf{3})$ was given by $H_R H_R / M_5$ as (70). Otherwise renormalization group equation (RGE) does not converge as we will show in the next paragraph. Thus in contrast with renormalizable $SO(10)$ where $\overline{\mathbf{126}}$ takes part in all mass matrices (See (7)), there appear independent $M_{15,2,2}$, $M_{10,1,3}$, $M_{10,3,1}$ in (68) and low energy data fitting is more easily satisfied.

In the following conveniences, let us introduce the following notations:

$$\begin{aligned} H_1 &= (\mathbf{1}, \mathbf{2}, \mathbf{2})_H, \quad H'_1 = (\mathbf{1}, \mathbf{2}, \mathbf{2})'_H, \\ H_6 &= (\mathbf{6}, \mathbf{1}, \mathbf{1})_H, \quad H_{15} = (\mathbf{15}, \mathbf{1}, \mathbf{1})_H, \\ H_L &= (\mathbf{4}, \mathbf{2}, \mathbf{1})_H, \quad \overline{H}_L = (\overline{\mathbf{4}}, \mathbf{2}, \mathbf{1})_H, \\ H_R &= (\mathbf{4}, \mathbf{1}, \mathbf{2})_H, \quad \overline{H}_R = (\overline{\mathbf{4}}, \mathbf{1}, \mathbf{2})_H. \end{aligned} \quad (69)$$

Superpotential relevant for fermion masses is given by⁷

$$\begin{aligned} W_Y &= Y_1^{ij} F_{Li} \overline{F_{Rj}} H_1 + \frac{Y_{15}^{ij}}{M_5} F_{Li} \overline{F_{Rj}} (H'_1 H_{15}) \\ &+ \frac{Y_R^{ij}}{M_5} \overline{F_{Ri}} \overline{F_{Rj}} (H_R H_R) + \frac{Y_L^{ij}}{M_5} F_{Li} F_{Lj} (\overline{H}_L \overline{H}_L), \end{aligned} \quad (70)$$

⁷ For simplicity, we have introduced only minimal terms necessary for reproducing observed fermion mass matrices.

On the other hand, the $(\overline{\mathbf{10}}, \mathbf{3}, \mathbf{1})$ and $(\mathbf{10}, \mathbf{1}, \mathbf{3})$ in ϕ_{126} were responsible for the left- and the right- handed Majorana neutrino masses and the same ϕ_{126} committed in the Yukawa coupling. This gave the severe constraints in the minimal SO(10) GUT. However in the present case Y_R^{ij} are coming from $\overline{H_R H_R}$ and are quite independent on the other Dirac Yukawa couplings and we have no problem in low energy data fitting including the neutrino oscillation data. Hereafter we consider type I seesaw since it is sufficient for the neutrino data fitting as indicated just above, and neglect the last term in (70). We introduce Higgs superpotential invariant under the PS symmetry such as

$$\begin{aligned} W = & \frac{m_1}{2} H_1^2 + \frac{m'_1}{2} H_1'^2 + m_{15} \text{tr} [H_{15}^2] + m_4 (\overline{H_L} H_L + \overline{H_R} H_R) \\ & + (H_L \overline{H_R} + \overline{H_L} H_R) (\lambda_1 H_1 + \lambda'_1 H'_1) + \lambda_{15} (\overline{H_R} H_R + \overline{H_L} H_L) H_{15} \\ & + \lambda \text{tr} [H_{15}^3] + \lambda_6 (H_L^2 + \overline{H_L}^2 + H_R^2 + \overline{H_R}^2) H_6. \end{aligned} \quad (71)$$

Parameterizing $\langle H_{15} \rangle = \frac{v_{15}}{2\sqrt{6}} \text{diag}(1, 1, 1, -3)$, SUSY vacuum conditions from Eq. (71) and the D-terms are satisfied by solutions,

$$v_{15} = \frac{2\sqrt{6}}{3\lambda_{15}} m_4, \quad \langle H_R \rangle = \langle \overline{H_R} \rangle = \sqrt{\frac{8m_4}{3\lambda_{15}^2} \left(m_{15} - \frac{\lambda}{\lambda_{15}} m_4 \right)} \equiv v_{PS} \quad (72)$$

and others are zero, by which the PS gauge symmetry is broken down to the SM gauge symmetry. We choose the parameters so as to be $v_{15} \simeq \langle H_R \rangle = \langle \overline{H_R} \rangle$. Note that the last term in Eq. (71) is necessary to make all color triplets in H_R and $\overline{H_R}$ heavy.

Weak Higgs doublet mass matrix is given by

$$\begin{pmatrix} H_1 & H'_1 & H_L \end{pmatrix} \begin{pmatrix} m_1 & 0 & \lambda_1 \langle H_R \rangle \\ 0 & m'_1 & \lambda'_1 \langle H_R \rangle \\ \lambda_1 \langle \overline{H_R} \rangle & \lambda'_1 \langle \overline{H_R} \rangle & m_4 \end{pmatrix} \begin{pmatrix} H_1 \\ H'_1 \\ \overline{H_L} \end{pmatrix}. \quad (73)$$

In order to realize the MSSM at low energy, only one pair of Higgs doublets out of the above tree pairs should be light, while others have mass of the PS symmetry breaking scale. This doublet-doublet Higgs mass splitting requires the fine tuning of parameters to satisfy

$$\det M = m_1 m'_1 m_4 - (m_1 \lambda_1'^2 + m'_1 \lambda_1^2) v_{PS}^2 = 0. \quad (74)$$

RGE. In our set up, the evolution of gauge coupling has three stages, G_{321} (SM+MSSM), G_{422} (the PS) and M_c ($\equiv (1/R)$) stages. In the G_{321} stage, the beta functions b_i are defined by

$$\frac{1}{\alpha_i(\mu)} = \frac{1}{\alpha_i(M)} + \frac{1}{2\pi} b_i \ln \left(\frac{M}{\mu} \right). \quad (i = 3, 2, 1) \quad (75)$$

b_i at G_{321} are

$$b_3 = -7, \quad b_2 = -19/6, \quad b_1 = 41/10 \quad (76)$$

at $M_{SUSY} > \mu > M = M_Z$ and

$$b_3 = -3, \quad b_2 = 1, \quad b_1 = 33/5 \quad (77)$$

at $M_c > \mu > M = M_{SUSY}$ The PS symmetry is recovered at $\mu = v_{PS}$. However, we assume $v_{PS} = M_c$ for simplicity, and the matching condition holds

$$\begin{aligned} \alpha_3^{-1}(M_c) &= \alpha_4^{-1}(M_c) \\ \alpha_2^{-1}(M_c) &= \alpha_{2L}^{-1}(M_c) \\ \alpha_1^{-1}(M_c) &= [2\alpha_4^{-1}(M_c) + 3\alpha_{2R}^{-1}]/5 \end{aligned} \quad (78)$$

at M_c . In the PS stage $\mu > M_c$, the threshold corrections Δ_i due to KK mode in the bulk are added,

$$\frac{1}{\alpha_i(\mu)} = \frac{1}{\alpha_i(M_c)} + \frac{1}{2\pi} b_i \ln \left(\frac{M_c}{\mu} \right) + \Delta_i. \quad (i = 4, 2_L, 2_R) \quad (79)$$

The beta functions of the PS gauge coupling constants are from the contents of Tables 3 and 4

$$b_4 = 3, \quad b_{2L} = b_{2R} = 6. \quad (80)$$

Δ_i are

$$\Delta_i = \frac{1}{2\pi} b_i^{even} \sum_{n=0}^{N_i} \theta(\mu - (2n+2)M_c) \ln \frac{(2n+2)M_c}{\mu} + \frac{1}{2\pi} b_i^{odd} \sum_{n=0}^{N_i} \theta(\mu - (2n+1)M_c) \ln \frac{(2n+1)M_c}{\mu} \quad (81)$$

with

$$\begin{aligned} b_i^{even} &= (-8, -4, -4) \\ b_i^{odd} &= (-8, -12, -12) \end{aligned} \quad (82)$$

under G_{422} .

In Fig.3 we depict the gauge coupling unification for left-right symmetric case. For simplicity we first assumed $M_c = v_{PS}$. The point v_{PS} is given by $\alpha_{2L}^{-1}(v_{PS}) = \alpha_2^{-1}(v_{PS}) = \alpha_{2R}^{-1}(v_{PS}) = (5\alpha_1^{-1} - 2\alpha_3)/3|_{v_{PS}}$. M_* is the scale of gauge coupling unification. The result is

$$v_{PS} = M_c = 1.19 \times 10^{16} \text{GeV}, \quad M_{GUT} = 4.61 \times 10^{17} \text{GeV} \quad (83)$$

In SO(10) model in 4D, $v_{PS} = O(10^{14} \text{GeV})$ was preferable for neutrino masses. However, it is not the case for the theories of SO(10) in 5D or the PS model in 4D. As is easily seen from (70),

$$M_R \sim Y_R v_{PS}^2 / M_5 \sim 0.1 Y_R v_{PS} \quad (84)$$

So if we assume $Y_R = 0.1$ we obtain the reasonable value of light neutrino mass.




FIGURE 3. Gauge coupling unification in the left-right symmetric case. Each line from top to bottom corresponds to g_3 , g_2 and g_1 for $\mu < M_c$, while $g_3 = g_4$ and $g_2 = g_{2R}$ for $\mu > M_c$.

For the case of no left-right symmetry, we can not fix v_{PS} as before. v_{PS} and M_c are determined from the requirement of gauge coupling unification.

gaugino mediation and gravitino problem. At $y = 0$ brane, $SO(10)$ remained invariant but supersymmetry is broken by the F-term of $SO(10)$ singlet S , $F_S \theta^2$. The interaction between S and bulk fields is

$$L = \delta(y) \int d^2\theta \lambda \frac{S}{M_5^2} \text{tr}[W_a W_a], \quad (85)$$

where λ is a dimensionless coupling constant. The bulk gaugino first obtains the mass

$$M_\lambda = \frac{\lambda F_S M_c}{M_5^2} \approx \frac{\lambda F_S M_5}{M_{PL}^2}, \quad (86)$$

where we have used $M_5^3/M_c \approx M_{PL}^2$ in the last equality, and M_c comes from the wave function normalization of the bulk gaugino.

In our scenario, M_5 is roughly one order of magnitude smaller than the (reduced) Planck mass (M_{PL}). As usual, we take $M_\lambda = 100\text{GeV} - 1\text{TeV}$. With this gaugino mass at high energy scale, SUSY breaking mass terms of sfermions are automatically generated through its RGE running and flavour blind citeGiudice. Comparing the gaugino mass to gravitino mass $m_{3/2} \approx F_S/M_{PL}$, a typical gaugino mass is smaller than the gravitino mass by a factor $\lambda M_5/M_{PL} \approx 0.1\lambda$. However, in this simple setup, it turns out that stau is the lightest superpartner (LSP), which is problematic in cosmology [62]. It has been found that when $M_c > M_{GUT}$, the RGE running in a unified theory pushes up stau mass and leads neutralino to be the LSP [63]. However, in our model, we cannot take such an arrangement, because M_c and M_{GUT} are fixed as $M_c < M_{GUT}$ to realize the gauge coupling unification. In order to avoid this problem, we need to extend the SUSY breaking sector. It is possible to introduce the gauge mediation [64] on the PS brane, in which gravitino is normally the LSP. In general, we can introduce the messenger sector on the brane at $y = 0$. This setup is basically the same as in Ref. [65], where the gauge mediation was calculated in 5D with the messenger sector on one brane, sfermions on the other brane and gauge multiplets in the bulk. When the messenger scale is larger than the compactification scale ($M_{mess} > M_c$), the gaugino mass is given by the same formula as in 4D,

$$M_\lambda \simeq \frac{\alpha_{GUT}}{4\pi} \frac{F_S}{M_5}, \quad (87)$$

while sfermion masses are roughly given by

$$\tilde{m}^2 \simeq M_\lambda^2 \left(\frac{M_c}{M_5} \right)^2. \quad (88)$$

The sfermion mass squared is suppressed relative to the gaugino mass by a geometric factor M_c/M_{mess} , at the messenger scale. At low energy, sfermion masses comparable to the gaugino mass are generated through the RGE running. In this setup, we find

$$\frac{m_{3/2}}{M_\lambda} \simeq \frac{M_{mess}}{\frac{\alpha_{GUT}}{4\pi} M_{PL}} > 10 \quad (89)$$

for $M_{mess} \geq M_c$. Thus, in order to have gravitino the LSP, the messenger scale should be smaller than the compactification scale. In this case, soft mass formulas are reduced into the usual four dimensional ones in the gauge mediation scenario.

Conclusion

We have reviewed the present status of renormalizable minimal $SO(10)$ GUT and tried to solve the problems by extending the theory in 4D into 5D. In the case of warped GUT, it not only solves the problems but also cures the blowup problem of the unified coupling after GUT, which had been thought as the fatal deficit of renormalizable $SO(10)$ GUT from the side of perturbative $SO(10)$ GUT group. Other problematic point of high dimension Higgs, if any, is that the complexity of intermediate energy scales (210 has three SM singlets (22)-(24)) is transformed to the

variety of Higgs profile in warped 5D, or is relaxed by using the freedom of the PS invariance in the case of orbifold GUT and gauge coupling unification is recovered. SO(10) GUT in 5D may also give the device of SUSY breaking mechanism, which was given by hand in 4D. Probably the more elaborate theory may be the warped orbifold GUT. The final theory may lead to 10D superstring or heterotic string theories [4][5][53] but 5D GUT may give the essential picture towards it. We should offer more elaborate arguments why an extra dimension is needed. Maldacena conjecture [66] is very suggestive for it.

Lastly I want to emphasize that the renormalizable minimal SO(10) GUT in 4D still remains valid as the essential part of the future complete theory. It is indeed still premature but "Don't throw the baby out with the bath water."

ACKNOWLEDGMENTS

As the chair of organizers, the author is grateful to Yukawa Institute for Theoretical Physics, Kyoto University and Ritsumeikan University for their financial supports. He also thanks all the organizers and participants at this Workshop.

REFERENCES

1. T.Fukuyama, AIP Conference Proceedings 1015 Published July 2008(Editors T.Fukuyama and T.Kikuchi): Some minor errors in section orbifold GUT are corrected.
2. H. Georgi and S.L. Glashow, Phys. Rev. Lett. **32** 438 (1974)
3. H. Murayama, How the neutrino physics requires the theory beyond SM is comprehensively discussed in this Proceedings.
4. J.E.Kim, this Proceeding
5. T. Kobayashi, S. Raby, R-J Zhang, Nucl.Phys. **B704** 3-55 (2005)
6. N. Maekawa, this Proceeding
7. S.King, This Proceedings.
8. T. Yanagida, in *Proceedings of the workshop on the Unified Theory and Baryon Number in the Universe*, edited by O. Sawada and A. Sugamoto (KEK, Tsukuba, 1979); M. Gell-Mann, P. Ramond, and R. Slansky, in *Supergravity*, edited by D. Freedman and P. van Nieuwenhuizen (north-Holland, Amsterdam 1979); R. N. Mohapatra and G. Senjanović, Phys. Rev. Lett. **44**, 912 (1980).
9. See also
L. J. Hall, R. Rattazzi and U. Sarid, Phys. Rev. D **50**, 7048 (1994) [arXiv:hep-ph/9306309]; G. Anderson, S. Raby, S. Dimopoulos, L. J. Hall and G. D. Starkman, Phys. Rev. D **49**, 3660 (1994) [arXiv:hep-ph/9308333]; L. J. Hall and S. Raby, Phys. Rev. D **51**, 6524 (1995) [arXiv:hep-ph/9501298]; R. Rattazzi and U. Sarid, Phys. Rev. D **53**, 1553 (1996) [arXiv:hep-ph/9505428]; C. H. Albright, K. S. Babu and S. M. Barr, Phys. Rev. Lett. **81**, 1167 (1998) [arXiv:hep-ph/9802314]; K. S. Babu, J. C. Pati and F. Wilczek, Nucl. Phys. B **566**, 33 (2000) [arXiv:hep-ph/9812538]; C. H. Albright and S. M. Barr, Phys. Rev. Lett. **85**, 244 (2000) [arXiv:hep-ph/0002155]; T. Blazek, R. Dermisek and S. Raby, Phys. Rev. D **65**, 115004 (2002) [arXiv:hep-ph/0201081]; D. Chang, T. Fukuyama, Y. Y. Keum, T. Kikuchi and N. Okada, Phys. Rev. D **71**, 095002 (2005) [arXiv:hep-ph/0412011].
10. K. S. Babu and R. N. Mohapatra, Phys. Rev. Lett. **70**, 2845 (1993) [arXiv:hep-ph/9209215].
11. K. Matsuda, Y. Koide and T. Fukuyama, Phys. Rev. D **64**, 053015 (2001) [arXiv:hep-ph/0010026]; K. Matsuda, Y. Koide, T. Fukuyama and H. Nishiura, Phys. Rev. D **65**, 033008 (2002) [Erratum-ibid. D **65**, 079904 (2002)] [arXiv:hep-ph/0108202].
12. T. Fukuyama and N. Okada, JHEP **0211**, 011 (2002) [arXiv:hep-ph/0205066].
13. M. Fukugita and T. Yanagida, Phys. Lett. B **174**, 45 (1986); for a recent review, see, W. Buchmuller, R. D. Peccei and T. Yanagida, Ann. Rev. Nucl. Part. Sci. **55**, 311 (2005) [arXiv:hep-ph/0502169].
14. T. Fukuyama, T. Kikuchi, N. Okada, Phys. Rev. D **68**, 033012 (2003) [arXiv:hep-ph/0304190].
See for LFV of different models, C.H. Albright, This Proceeding.
15. For a recent review, see, for example, J. Hisano, Nucl. Phys. Proc. Suppl. **111**, 178 (2002) [arXiv:hep-ph/0204100], and references therein.
16. J. Hisano, T. Moroi, K. Tobe and M. Yamaguchi, Phys. Rev. D **53**, 2442 (1996) [arXiv:hep-ph/9510309].
17. Y. Okada, K. i. Okumura and Y. Shimizu, Phys. Rev. D **61**, 094001 (2000) [arXiv:hep-ph/9906446].
18. C. L. Bennett *et al.* [WMAP Collaboration], Astrophys. J. Suppl. **148**, 1 (2003) [arXiv:astro-ph/0302207]; D. N. Spergel *et al.* [WMAP Collaboration], Astrophys. J. Suppl. **148**, 175 (2003) [arXiv:astro-ph/0302209].
19. J. R. Ellis, K. A. Olive, Y. Santoso and V. C. Spanos, Phys. Lett. B **565**, 176 (2003) [arXiv:hep-ph/0303043]; A. B. Lahanas and D. V. Nanopoulos, Phys. Lett. B **568**, 55 (2003) [arXiv:hep-ph/0303130].
20. T. Fukuyama, A. Ilakovac, T. Kikuchi and S. Meljanac, Nucl. Phys. Proc. Suppl. **144**, 143 (2005) [arXiv:hep-ph/0411282]; T. Fukuyama, A. Ilakovac and T. Kikuchi, arXiv:hep-ph/0506295.
21. K. Eguchi *et al.* [KamLAND Collaboration], Phys. Rev. Lett. **90**, 021802 (2003) [arXiv:hep-ex/0212021].
22. B. Bajc, G. Senjanović and F. Vissani, Phys. Rev. Lett. **90**, 051802 (2003) [arXiv:hep-ph/0210207]; H. S. Goh, R. N. Mohapatra and S. P. Ng, Phys. Lett. B **570**, 215 (2003) [arXiv:hep-ph/0303055].

23. H. S. Goh, R. N. Mohapatra and S. P. Ng, Phys. Rev. D **68**, 115008 (2003) [arXiv:hep-ph/0308197]; B. Dutta, Y. Mimura and R. N. Mohapatra, Phys. Rev. D **69**, 115014 (2004) [arXiv:hep-ph/0402113]; *ibid.*, Phys. Lett. B **603**, 35 (2004) [arXiv:hep-ph/0406262]; S. Bertolini, M. Frigerio and M. Malinsky, Phys. Rev. D **70**, 095002 (2004) [arXiv:hep-ph/0406117]; S. Bertolini and M. Malinsky, Phys. Rev. D **72**, 055021 (2005) [arXiv:hep-ph/0504241].
24. S. Bertolini and M. Malinsky, Phys. Rev. D **72**, 055021 (2005) [arXiv:hep-ph/0504241].
25. K. S. Babu and C. Macesanu, Phys. Rev. D **72**, 115003 (2005) [arXiv:hep-ph/0505200].
26. For the early work of threshold correction, see, D. Chang, R. N. Mohapatra and M. K. Parida, Phys. Rev. D **30**, 1052 (1984).
27. T. Fukuyama, A. Ilakovac, T. Kikuchi, S. Meljanac and N. Okada, J. Math. Phys. **46** 033505 (2005)
28. T. E. Clark, T. K. Kuo and N. Nakagawa, Phys. Lett. B **115**, 26 (1982); C. S. Aulakh and R. N. Mohapatra, Phys. Rev. D **28**, 217 (1983).
29. D. G. Lee, Phys. Rev. D **49**, 1417 (1994).
30. C. S. Aulakh, B. Bajc, A. Melfo, G. Senjanović and F. Vissani, Phys. Lett. B **588**, 196 (2004) [arXiv:hep-ph/0306242].
31. X. G. He and S. Meljanac, Phys. Rev. D **41** (1990) 1620.
32. J. Sato, Phys. Rev. D **53**, 3884 (1996) [arXiv:hep-ph/9508269].
33. T. Fukuyama, A. Ilakovac, T. Kikuchi, S. Meljanac and N. Okada, Eur. Phys. J. C **42**, 191 (2005) [arXiv:hep-ph/0401213]; *ibid.*, J. Math. Phys. **46**, 033505 (2005) [arXiv:hep-ph/0405300]; *ibid.*, Phys. Rev. D **72**, 051701 (2005) [arXiv:hep-ph/0412348].
34. B. Bajc, A. Melfo, G. Senjanović and F. Vissani, Phys. Rev. D **70**, 035007 (2004) [arXiv:hep-ph/0402122].
35. C. S. Aulakh and A. Girdhar, Nucl. Phys. B **711**, 275 (2005) [arXiv:hep-ph/0405074].
36. S. Bertolini, T. Schwetz and M. Malinsky, Phys. Rev. D **73**, 115012 (2006) [arXiv:hep-ph/0605006].
37. Y. Mimura, This Proceeding.
38. T. Fukuyama, T. Kikuchi and N. Okada, arXiv:hep-ph/0702048.
39. T. Fukuyama, A. Ilakovac, T. Kikuchi, S. Meljanac and N. Okada, JHEP **0409** 052 (2004) [arXiv:hep-ph/0406068]
Discrete symmetry for avoiding fast proton decay was discussed by
R.N. Mohapatra, in this Proceedings.
40. M. Luty, 2004 TASI Lectures on Supersymmetry Breaking, arXiv:hep-ph/0509029
41. L. Randall and R. Sundrum, Phys. Rev. Lett. **83**, 3370 (1999) [arXiv:hep-ph/9905221].
42. Y. Nomura, D. Poland and B. Tweedie, JHEP **0612**, 002 (2006) [arXiv:hep-ph/0605014].
43. M. A. Luty and R. Sundrum, Phys. Rev. D **64**, 065012 (2001) [arXiv:hep-th/0012158].
44. J. Bagger, D. Nemeschansky and R. J. Zhang, JHEP **0108**, 057 (2001) [arXiv:hep-th/0012163].
45. D. Marti and A. Pomarol, Phys. Rev. D **64**, 105025 (2001) [arXiv:hep-th/0106256].
46. R. Kitano and T. j. Li, Phys. Rev. D **67**, 116004 (2003) [arXiv:hep-ph/0302073].
47. Y. Kawamura, Prog. Theor. Phys. **105**, 999 (2001) [arXiv:hep-ph/0012125].
48. G. Altarelli and F. Feruglio, Phys. Lett. B **511**, 257 (2001) [arXiv:hep-ph/0102301]; A. B. Kobakhidze, Phys. Lett. B **514** 131 (2001) [arXiv:hep-ph/0102323]
49. L. J. Hall and Y. Nomura, Phys. Rev. D **64**, 055003 (2001) [arXiv:hep-ph/0103125].
50. Y. Grossman and M. Neubert, Phys. Lett. B **474**, 361 (2000) [arXiv:hep-ph/9912408].
51. S. Chang, J. Hisano, H. Nakano, N. Okada and M. Yamaguchi, Phys. Rev. D **62**, 084025 (2000) [arXiv:hep-ph/9912498].
52. T. Gherghetta and A. Pomarol, Nucl. Phys. B **586**, 141 (2000) [arXiv:hep-ph/0003129].
53. K.-S. Choi and J.E. Kim, Quarks and Leptons from Orbifolded Superstring, Springer 2006;
54. T. Asaka, W. Buchmuler and L. Covi, Phys. Lett. **B523** 199 (2001) L.J. Hall, Y. Nomura, T. Okui and D. Smith, Phys. Rev. **D65** 035008 (2002) L.J. Hall and Y. Nomura, Phys. Lett. **703** 217 (2004) B. Kyae, C.A. Lee and Q. Shafi, Nucl. Phys. **B683** 105 (2004)
55. Y. Kawamura, Prog. Theor. Phys. **105**, 999 (2001) [arXiv:hep-ph/0012125].
56. L. Hall and Y. Nomura, Phys. Rev. D **64** 055003 (2001)
57. R. Dermisek and A. Mafi, Phys. Rev. D **65**, 055002 (2002) H.D. Kim and S. Raby, JHEP **0301** 056 (2003)
58. I. Dorsner, Phys. Rev. D **69** 056003 (2004)
59. K. Matsuda, T. Fukuyama and H. Nishiura Physical Review D **61**, 053001 (2000). [arXiv:hep-ph/9906433]
60. R. Mohapatra and R.E. Marshak, Phys. Rev. Lett. **44** 1316 (1980)
61. G.F. Giudice and A. Masiero, Phys. Lett. B **377** 480 (1988)
62. As for the gravitino problem, M. Kawasaki, This Proceedings. See also
M. Y. Khlopov and A. D. Linde, Phys. Lett. B **138**, 265 (1984); J. R. Ellis, J. E. Kim and D. V. Nanopoulos, Phys. Lett. B **145**, 181 (1984); M. Kawasaki, K. Kohri, and T. Moroi, Phys. Rev. D **71** 083502 (2005); K. Kohri, M. Yamaguchi, and J. Yokoyama, Phys. Rev. D **72** 083510 (2005) [arXiv:hep-ph/0502211]
On the broad confrontation with Cosmology; Q. Shafi, This Proceedings and references therein.
63. M. Schmaltz and W. Skiba, Phys. Rev. D **62**, 095004 (2000).
64. For a review, see G. F. Giudice and R. Rattazzi, Phys. Rept. **322**, 419 (1999), and references therein.
65. E. A. Mirabelli and M. E. Peskin, Phys. Rev. D **58**, 065002 (1998).
66. J. Maldacena, Adv. Theor. Math. Phys. **2** 231 (1998)